

Characteristics and Outcomes of Women Veterans Undergoing Cardiac Catheterization in the Veterans Affairs Healthcare System Insights from the VA CART Program

Melinda B. Davis, MD; Thomas M. Maddox, MD, MSc; Paula Langner, MS;
Mary E. Plomondon, PhD, MSPH; John S. Rumsfeld, MD, PhD; Claire S. Duvernoy, MD

Background—The number of women veterans is increasing, yet little is known about their cardiovascular risk factors, coronary anatomy, cardiac treatments, and outcomes after cardiac catheterization. Prior studies have shown that nonveteran women have more risk factors, receive less aggressive treatment, and have worse outcomes, despite having less obstructive coronary artery disease than men. Whether these differences exist among women veterans in the veterans affairs healthcare system is unknown.

Methods and Results—Data on 85 936 veterans (3181 women) undergoing initial cardiac catheterization between October 1, 2007, and September 30, 2012, were examined using the national veterans affairs Clinical Assessment Reporting and Tracking (CART) Program. Sex differences in demographics, indications, coronary anatomy, cardiac treatments, and outcomes were analyzed. Women veterans were younger (56.9 versus 63.0 years, $P < 0.0001$) with fewer traditional cardiovascular risk factors, but with more obesity, depression, and posttraumatic stress disorder than men. Women had lower rates of obstructive coronary artery disease than men (22.6% versus 53.3%). Rates of procedural complications were similar in both genders. Adjusted outcomes at 1 year showed women had lower mortality (hazard ratio, 0.74; confidence interval, 0.60–0.92) and less all-cause rehospitalization (hazard ratio, 0.87; confidence interval, 0.82–0.93), but no difference in rates of unplanned percutaneous coronary intervention.

Conclusions—Women veterans undergoing catheterization are younger, have more obesity, depression, and posttraumatic stress disorder, less obstructive coronary artery disease, and similar long-term outcomes, compared with men. These findings suggest a significant portion of women veterans may have chest pain not attributable to obstructive coronary artery disease. Further research into possible causes, such as endothelial dysfunction or concurrent psychological comorbidities, is needed. (*Circ Cardiovasc Qual Outcomes*. 2015;8:S39-S47. DOI: 10.1161/CIRCOUTCOMES.114.001613.)

Key Words: cardiac catheterization ■ coronary artery disease ■ outcomes research ■ sex ■ women

Coronary artery disease (CAD) remains the leading cause of death among women in the United States.¹ Prior research has demonstrated that women with CAD are typically older than men and have more risk factors, such as diabetes mellitus,² hypertension,^{3,4} hyperlipidemia,⁵ obesity,⁶ and congestive heart failure.^{4,7} Despite these higher rates of CAD risk factors, women have less obstructive CAD.^{4,8} Studies are conflicting regarding whether women are treated less aggressively than their male counterparts.^{9–15} Women tend to have more complications and worse outcomes after cardiac catheterization¹⁶ and are less likely to receive evidence-based medical therapy after discharge.¹⁷

Women who have served in the military may have an even higher risk of CAD than nonveteran women, given the overall worse health status and higher number of CAD risk factors in the veteran population,^{18–20} but little is known about the characteristics and treatment of women veterans suspected of having clinically significant CAD. The number of women seeking care at veterans affairs (VA) facilities has doubled over the past decade,²¹ providing a unique opportunity to understand cardiac care and outcomes among women veterans. By extension, these lessons could improve our understanding of the sex differences noted in nonveteran cardiac populations.

Received December 10, 2014; accepted January 23, 2015.

From the VA Ann Arbor Healthcare System, Ann Arbor, MI (M.B.D., C.S.D.); University of Michigan Healthcare System, Ann Arbor, MI (M.B.D., C.S.D.); VA Eastern Colorado Health Care System, Denver, CO (T.M.M., P.L., M.E.P., J.S.R.); and University of Colorado School of Medicine, Denver, CO (T.M.M., J.S.R.).

This article was handled independently by Robert W. Yeh, MD, MSc, as a Guest Editor. The editors had no role in the evaluation of the manuscript or in the decision about its acceptance.

Correspondence to Claire Duvernoy, MD, Cardiology Section, Medicine Service, VA Ann Arbor Healthcare System, 2215 Fuller Rd, Box 111a, Ann Arbor, MI 48105. E-mail duvernoy@umich.edu

© 2015 American Heart Association, Inc.

Circ Cardiovasc Qual Outcomes is available at <http://circoutcomes.ahajournals.org>

DOI: 10.1161/CIRCOUTCOMES.114.001613

WHAT IS KNOWN

- Despite having less obstructive coronary artery disease than men, women have higher rates of cardiovascular risk factors and worse cardiovascular outcomes.
- It is not known whether these characteristics apply to women veterans undergoing cardiac catheterization in the VA healthcare system.

WHAT THE STUDY ADDS

- This national study demonstrates that women veterans undergoing cardiac catheterization in the VA system have higher rates of nonobstructive coronary artery disease or normal coronary arteries than men, even when presenting with acute coronary syndromes.
- Women veterans have a higher burden of obesity, depression, and posttraumatic stress disorder, fewer traditional cardiovascular risk factors, and similar 1-year outcomes compared with men.
- Chest pain in the absence of obstructive coronary disease is common in women veterans and may represent a complex interplay of psychological stressors and somatic disease, but further research is needed.

Using data from the national VA Clinical Assessment, Reporting, and Tracking (CART) Program, we sought to evaluate sex differences among veterans undergoing initial diagnostic catheterization in the VA. The specific aims of this study were to determine whether there were sex differences in (1) clinical characteristics and comorbidities, (2) coronary anatomy and treatment, and (3) procedural complications and long-term outcomes after diagnostic catheterization.

Methods

Launched in 2005, the VA CART Program is a national VA clinical quality program for coronary procedures conducted in all VA cardiac catheterization laboratories nationwide.²² A key feature of the CART Program is a clinical software application designed to collect data on catheterization laboratory procedures in a standardized fashion. The software is embedded in the VA electronic health record and allows providers to enter patient and procedural information for all cardiac catheterizations and percutaneous coronary interventions (PCI) as part of routine clinical workflow. Once data entry is complete, the data elements automatically populate a clinical note for the patient electronic health record. These data also populate a clinical data repository that supports the quality assessment, quality improvement, and clinical research missions of the CART Program.

The CART software uses standardized definitions and features, such as pull-down menus, to ensure uniformity of data entry by different providers and in different catheterization laboratories. Core data elements conform to the definitions and standards of the American College of Cardiology's National Cardiovascular Data Registry.²³ Quality checks of the data are periodically conducted, and it has been shown to be complete, timely, and accurate.²⁴ CART data are combined with longitudinal data in the VA patient electronic health record, including vital status, inpatient hospitalizations, clinic visits, pharmacy prescriptions and refills, and laboratory results. The data are also merged with VA claims data for veterans' care for

hospitalizations that occur outside the VA system. The Colorado Multiple Institutional Review Board approved this study.

Study Cohort

This study included data from all men and women veterans undergoing initial diagnostic catheterization between October 1, 2007, and September 30, 2012, at any of the 77 VA catheterization laboratories. Patients <18 years of age were excluded. All patients with a prior catheterization, PCI, or coronary artery bypass grafting (CABG) were excluded to evaluate the risk factor profile and subsequent care after the initial catheterization.

Exposure

Patient sex was determined from the CART database and VA patient treatment files.

Outcomes

Both peri-procedural and 1-year outcomes were assessed. Peri-procedural outcomes included in-laboratory complications and bleeding complications requiring transfusion. One-year outcomes included all-cause death, all-cause rehospitalization, rehospitalization for myocardial infarction (MI), unplanned catheterization, unplanned PCI, and unplanned CABG. In-laboratory complications were defined as access-site hematoma, dysrhythmia, hives, stroke, emergent PCI, acute respiratory distress, acute pulmonary edema, death, periprocedural MI, acute cardiogenic shock, limb ischemia, anaphylactic shock, retroperitoneal hematoma, emergent CABG, or new cardiac tamponade.

Covariates

Patient demographic, clinical comorbidity, catheterization indication, coronary anatomy, and postprocedural treatment information were collected from the CART and VA patient data files. Demographic information included age, race, and Hispanic ethnicity. Race was defined as white, black or African American, or other, which included American Indian or Alaska Native, Asian, Native Hawaiian, or other Pacific Islander. Racial and ethnic classifications were based on patient self-report, and the methodology for racial classification has been described elsewhere.²⁵ Clinical comorbidity information included diabetes mellitus, hyperlipidemia, hypertension, chronic kidney disease, tobacco use, congestive heart failure, cerebrovascular disease, peripheral arterial disease, obesity, chronic obstructive pulmonary disease, posttraumatic stress disorder (PTSD), depression, sleep apnea, procedural indication, and Framingham risk. Obesity was defined as a body mass index ≥ 30 . Framingham risk was calculated using methods previously described²⁶ and was categorized by the 10-year predicted coronary heart disease risk as low (<10%), intermediate (10% to 20%), or high (>20%). Indications for cardiac catheterization were collected from CART data and categorized into acute coronary syndrome (ACS), elective, or other. Elective indications included chest pain, stable angina, and positive functional study. Positive functional study was defined as any cardiac stress test that was suggestive of ischemia. Other indications for cardiac catheterization included valvular heart disease, cardiomyopathy, ischemic heart disease, cardiac transplant, cardiac tamponade, cardiogenic shock, coronary heart disease, heart failure, dysrhythmia, research study, and unknown.

Coronary anatomy was classified into obstructive CAD, nonobstructive CAD, or normal coronary arteries. In addition, the extent of CAD was classified into 1, 2, or 3-vessel distribution. In line with prior studies,²⁵ obstructive CAD was defined as any coronary stenosis >70% in any epicardial artery or 50% in the left main coronary artery. Nonobstructive CAD refers to coronary stenosis that is $\geq 20\%$, but <70%, in any epicardial artery or $\geq 20\%$, but <50%, in the left main coronary artery. Normal coronary anatomy was defined as <20% stenosis in all coronary arteries. We then categorized patients as having single, double, or triple-vessel or left main obstructive disease. Vessel

Table 1. Demographics and Clinical Characteristics

	Women (N=3181)	Men (N=82755)	P Value
Demographics			
Age, median (IQR)	56.9 (51.1–62.5)	63 (58.3–69)	<0.0001
Hispanic, % (n)	6.8 (216)	7.9 (6508)	0.0269
Race: White, % (n)	63.9 (2033)	73.4 (60756)	<0.0001
Race: Black, % (n)	25.9 (824)	17.4 (14394)	<0.0001
Race: Other, % (n)	10.2 (324)	9.2 (7605)	0.0569
Clinical Comorbidities			
Obese, % (n)	57.2 (1794)	46.6 (38149)	<0.0001
BMI, median (IQR)	31.2 (26.7–36.6)	29.5 (26–33.8)	<0.0001
Hypertension, % (n)	75.7 (2409)	82.7 (68469)	<0.0001
Hyperlipidemia, % (n)	71.2 (2265)	77.5 (64117)	<0.0001
Diabetes mellitus, % (n)	32.7 (1041)	39.4 (32644)	<0.0001
Depression, % (n)	55.3 (1758)	31.4 (26002)	<0.0001
Congestive heart failure, % (n)	13.4 (426)	20 (16551)	<0.0001
Cerebrovascular disease, % (n)	10.2 (323)	12.6 (10394)	<0.0001
Framingham risk: high, % (n)	2.9 (92)	24.6 (20337)	<0.0001
Framingham risk: medium, % (n)	38.6 (1229)	54.6 (45173)	<0.0001
Framingham risk: low, % (n)	58.5 (1860)	20.8 (17245)	<0.0001
Tobacco use, % (n)	48.9 (1556)	57.7 (47760)	<0.0001
Prior myocardial infarction (MI), % (n)	4.8 (152)	7.8 (6438)	<0.0001
COPD, % (n)	17.5 (557)	20 (16545)	0.0006
Chronic kidney disease, % (n)	6.6 (210)	14.2 (11714)	<0.0001
Family history of CAD, % (n)	1.3 (40)	0.5 (431)	<0.0001
Peripheral vascular disease, % (n)	6.9 (220)	14.9 (12309)	<0.0001
PTSD, % (n)	20.1 (638)	16.3 (13481)	<0.0001
Sleep apnea, % (n)	15.3 (488)	16.8 (13870)	0.0352
Pre-catheterization stress test			
Stress test \leq 30 days prior, % (n)	30.4 (966)	28.1 (23233)	0.0048

BMI indicates body mass index; CAD, coronary artery disease; COPD, chronic obstructive pulmonary disease; IQR, interquartile range; and PTSD, posttraumatic stress disorder.

distribution was classified into the left anterior descending and its branches, the left circumflex artery and its branches, and the right coronary artery and its branches. Postprocedural treatments included PCI, CABG, or medical therapy. PCI that occurred on the same day as the initial catheterization or as part of a staged PCI were considered part of the treatment episode and not counted as an outcome. CABG that occurred within 30 days of the index catheterization, without an intervening unplanned catheterization or PCI, was considered a planned procedure. All other CABG procedures were considered unplanned. Medications at the time of discharge after cardiac catheterization were analyzed, including β -blockers, statin therapy, antiplatelet agents (clopidogrel or prasugrel), long-acting nitrates, and calcium-channel blockers.

Statistical Analyses

Patient demographics and clinical characteristics, procedural indications, presence of coronary stenoses, pharmacological treatment, and revascularization procedures were collected and compared by sex. The cohort was further stratified and analyzed by indication (ACS, elective, or other). Continuous data were compared using Wilcoxon nonparametric tests and categorical data using Chi-square or Fisher's exact tests, as appropriate.

Unadjusted rates of peri-procedural and 1-year outcomes were calculated and compared using Kaplan–Meier curves. To evaluate adjusted differences in 1-year outcomes for men and women after initial cardiac catheterization, a Cox proportional hazards model was fitted for each of 6 outcomes, adjusted using the covariates listed earlier and including an interaction between sex and indication for catheterization to assess sex differences by indication category. Spline terms were included in the models to account for the nonlinearity of age. For all outcomes, patients were censored on death or at 1 year if they did not experience an event. Crude and adjusted estimates of the hazard ratios for women compared with men were calculated. Data were complete for all covariates with the exception of obesity, which resulted in exclusion of 920 (1.1%) patients. These patients were excluded from both adjusted and unadjusted models. Adjustment for the peri-procedural outcomes was not performed because of low numbers of events. Sex differences in clinical characteristics were also assessed by stratifying the cohort by race (white, black, and other).

Statistical analyses were performed using SAS software (version 9.4, SAS Institute, Cary, NC) and R 3.1.1 (R Core Team (2014)). All analyses were performed at the CART Coordinating Center, housed at the VA Eastern Colorado Healthcare System.

Results

Clinical Characteristics

A total of 82 755 males (96%) and 3181 (4%) women met criteria for inclusion in this analysis. The median age for women was 56.9 years and for men was 63.0 years ($P<0.0001$). Baseline demographics for the entire population revealed that women had fewer traditional cardiovascular risk factors than men, including hypertension, hyperlipidemia, diabetes mellitus, and tobacco use (Table 1). Notably, women veterans had higher rates of obesity, depression, and PTSD than men veterans (Table 1). The same findings were apparent when analyzed by race. Women in each racial group (white, black, and other) were younger, had lower rates of traditional cardiovascular risk factors (hypertension, hyperlipidemia, diabetes mellitus, and tobacco use), and higher rates of obesity, depression, and PTSD than their male counterparts.

Table 2 shows the demographics and clinical characteristics of veterans by indication for catheterization (ACS, elective, and other). Regardless of catheterization indication, women had lower rates of hypertension, hyperlipidemia, diabetes mellitus, and tobacco use than men. In each category, women had more obesity, depression, and PTSD than men. Among the entire cohort, women more commonly presented with chest pain as an indication for cardiac catheterization than men (52.1%, $n=1656$ versus 39.3%, $n=32549$, $P<0.0001$). Although stress testing was performed equally often in men and women, irrespective of indication for catheterization (Table 2), a relatively low percentage of patients presented for catheterization with the indication of positive functional study, and women were somewhat less likely to present with this indication than men (13.3%, $n=423$ versus 14.8%, $n=12212$, $P=0.023$).

Coronary Anatomy and Treatment

Women were found to have normal coronary arteries at the time of catheterization significantly more often than men (45.9% versus 17%, $P<0.0001$; Table 3). Rates of nonobstructive CAD were similar but women had less single and multivessel obstructive disease. Rates of treatment with PCI and medical therapy were lower in women (Table 3). Analysis

Table 2. Demographics and Clinical Characteristics by Indication

	ACS			Elective			Other		
	Women (N=401)	Men (N=14 052)	P Value	Women (N=2132)	Men (N=46 309)	P Value	Women(N=648)	Men (N=22 394)	P Value
Demographics									
Age, median (IQR)	58.7 (52.4–65.4)	63.5 (58.6–71.6)	<0.0001	56.2 (50.6–61.6)	62.6 (57.8–67.5)	<0.0001	58 (51.6–65)	63.8 (59.2–71.5)	<0.0001
Hispanic, % (n)	10.7 (43)	11.2 (1572)	0.7713	6.4 (136)	7.3 (3374)	0.1143	5.7 (37)	7 (1562)	0.2115
Race: White, % (n)	60.1 (241)	72.6 (10 199)	<0.0001	64.4 (1374)	73.7 (34 121)	<0.0001	64.5 (418)	73.4 (16 436)	<0.0001
Race: Black, % (n)	27.9 (112)	17.5 (2455)	<0.0001	25.9 (552)	17.1 (7896)	<0.0001	24.7 (160)	18.1 (4043)	<0.0001
Race: Other, % (n)	12 (48)	9.9 (1398)	0.1835	9.7 (206)	9.3 (4292)	0.5398	10.8 (70)	8.6 (1915)	0.0441
Clinical comorbidities									
Obese, % (n)	50 (188)	40.5 (5566)	0.0002	61.2 (1296)	50.1 (23 036)	<0.0001	48.1 (310)	43.1 (9547)	0.0116
BMI, median (IQR)	30 (25.3–34.8)	28.7 (25.2–32.7)	0.001	31.9 (27.6–37.2)	30 (26.6–34.3)	<0.0001	29.6 (25.2–35.5)	29 (25.3–33.4)	0.0065
Hypertension, % (n)	75.6 (303)	81.1 (11 391)	0.0057	75.8 (1617)	83.7 (38 750)	<0.0001	75.5 (489)	81.8 (18 328)	<0.0001
Hyperlipidemia, % (n)	66.1 (265)	74.2 (10 424)	0.0003	74 (1578)	81.1 (37 552)	<0.0001	65.1 (422)	72.1 (16 141)	0.0001
Diabetes mellitus, % (n)	31.7 (127)	38.9 (5473)	0.0032	33.6 (716)	40.3 (18 666)	<0.0001	30.6 (198)	38 (8505)	0.0001
Depression, % (n)	45.4 (182)	28.9 (4068)	<0.0001	58.7 (1251)	34.2 (15 844)	<0.0001	50.2 (325)	27.2 (6090)	<0.0001
Congestive heart failure, % (n)	12 (48)	15 (2106)	0.0944	6.5 (138)	11.9 (5522)	<0.0001	37 (240)	39.8 (8923)	0.1498
Cerebrovascular disease, % (n)	15 (60)	14.3 (2005)	0.6953	9.2 (197)	11.7 (5435)	0.0004	10.2 (66)	13.2 (2954)	0.0254
Framingham risk: high, % (n)	4.7 (19)	31 (4352)	<0.0001	2.5 (54)	23.8 (11 024)	<0.0001	2.9 (19)	22.2 (4961)	<0.0001
Framingham risk: medium, % (n)	47.9 (192)	53.4 (7497)	0.0304	37.9 (808)	54.7 (25 347)	<0.0001	35.3 (229)	55.1 (12 329)	<0.0001
Framingham risk: low, % (n)	47.4 (190)	15.7 (2203)	<0.0001	59.6 (1270)	21.5 (9938)	<0.0001	61.7 (400)	22.8 (5104)	<0.0001
Tobacco use, % (n)	54.1 (217)	58.5 (8226)	0.0763	49.5 (1055)	59.2 (27 398)	<0.0001	43.8 (284)	54.2 (12 136)	<0.0001
Prior myocardial infarction, % (n)	12.5 (50)	16 (2246)	0.0576	3.8 (81)	5.8 (2672)	0.0001	3.2 (21)	6.8 (1520)	0.0004
COPD, % (n)	18.5 (74)	19.3 (2715)	0.6643	15.9 (338)	18.8 (8699)	0.0007	22.4 (145)	22.9 (5131)	0.7489
Chronic kidney disease, % (n)	10 (40)	16.2 (2280)	0.0008	4.8 (103)	12 (5545)	<0.0001	10.3 (67)	17.4 (3889)	<0.0001
Family history of CAD, % (n)	1.7 (7)	0.4 (58)	0.0021	1.4 (29)	0.6 (298)	<0.0001	0.6 (4)	0.3 (75)	0.287
Peripheral vascular disease, % (n)	9.2 (37)	16.2 (2277)	0.0002	6 (128)	13.9 (6434)	<0.0001	8.5 (55)	16.1 (3598)	<0.0001
PTSD, % (n)	18 (72)	13.8 (1939)	0.0177	22.4 (477)	18.7 (8651)	<0.0001	13.7 (89)	12.9 (2891)	0.5373
Sleep apnea, % (n)	10 (40)	12.8 (1795)	0.0969	16.7 (356)	18.7 (8655)	0.0208	14.2 (92)	15.3 (3420)	0.4532
Pre-Catheterization stress test									
Stress test ≤30 days prior, % (n)	11.2 (45)	12.4 (1744)	0.4759	37.9 (809)	37.6 (17 435)	0.7825	17.3 (112)	18.1 (4054)	0.5933

BMI indicates body mass index; CAD, coronary artery disease; COPD, chronic obstructive pulmonary disease; IQR, interquartile range; and PTSD, posttraumatic stress disorder.

by race (white, black, and other) showed that women in all groups had less obstructive CAD and were less likely to be treated with PCI or medications (β-blockers, statins, clopidogrel or prasugrel, and long-acting nitrates).

When differentiated by indication for catheterization (Table 4), women continued to have higher rates of normal coronaries than men (19.2% versus 5.0%, $P<0.0001$, in the ACS group). Among patients with ACS, women had similar rates of single-vessel coronary disease, and higher rates of nonobstructive and normal coronaries, compared with men.

Women had less 2-vessel and 3-vessel or left main CAD than men, regardless of the indication for catheterization, and had less treatment with PCI. At discharge, women were less likely to receive treatment with β-blockers, statins, and antiplatelet agents (clopidogrel or prasugrel), even among patients presenting with ACS (Table 4).

Outcomes

In-laboratory complications were similar among all women and men veterans (1.5% versus 1.6%, $p=NS$). Women were

Table 3. Coronary Anatomy, Postprocedural Medications, and Unadjusted Outcomes

	Women (N=3181)	Men (N=82755)	P Value
Coronary anatomy and treatment			
3-Vessel/left main obstructive, % (n)	4.5 (142)	17.1 (14132)	<0.0001
2-Vessel Obstructive, % (n)	5.3 (170)	14.3 (11842)	<0.0001
1-Vessel Obstructive, % (n)	12.8 (407)	21.9 (18103)	<0.0001
Non-Obstructive, % (n)	28.7 (913)	27.2 (22482)	0.0563
Normal, % (n)	45.9 (1461)	17 (14108)	<0.0001
PCI-Treatment, % (n)	11.7 (372)	21.8 (18070)	<0.0001
Postprocedural Medications			
Beta-blocker, % (n)	55.5 (1764)	70.5 (58337)	<0.0001
Statin, % (n)	57.3 (1823)	71.2 (58907)	<0.0001
Clopidogrel, prasugrel, % (n)	16.8 (533)	32.1 (26540)	<0.0001
Long-acting Nitrate, % (n)	24.1 (768)	31.3 (25902)	<0.0001
Calcium channel blocker, % (n)	23 (733)	22.9 (18974)	0.8795
Procedural complications			
In-laboratory complications, % (n)	1.5 (49)	1.6 (1318)	0.8172
Transfusion within 30 days, % (n)	3.4 (109)	5.8 (4839)	<0.0001
Unadjusted 1-year outcomes			
Death, % (n)	2.7 (86)	5.6 (4668)	<0.0001
Rehospitalization All-Cause, % (n)	30.6 (973)	41.5 (34383)	<0.0001
Rehospitalization for MI (day 15-1yr), % (n)	0.8 (24)	1.2 (952)	0.0387
Repeat Catheterization, % (n)	7.5 (239)	11.8 (9776)	<0.0001
Unplanned PCI, % (n)	3 (95)	5.7 (4698)	<0.0001
Unplanned CABG, % (n)	1.3 (40)	5.1 (4227)	<0.0001

CABG indicates coronary artery bypass grafting; MI, myocardial infarction; and PCI, percutaneous coronary intervention.

less likely to have bleeding complications requiring transfusion. At 1 year, women had lower rates of death (2.7% versus 5.6%), unplanned PCI (3.0% versus 5.7%), and unplanned CABG (1.3% versus 5.1%; all $P<0.0001$) than men (Table 3). Women were less likely to be rehospitalized for all causes. Similar sex differences in long-term outcomes were demonstrated in proportional hazards models adjusted for the presence of obstructive disease as well as multiple factors. Women veterans had similar rates of unplanned PCI (hazard ratio [HR], 1.03; confidence interval [CI], 0.84–1.27; $P=0.78$) and rehospitalization for MI (HR, 1.19; CI, 0.79–1.80; $P=0.41$). Women were less likely than men to die (HR, 0.74; CI, 0.60–0.92; $P=0.007$), undergo unplanned CABG (HR, 0.58; CI, 0.42–0.79; $P<0.001$), or be rehospitalized for any cause (HR, 0.87; CI, 0.82–0.93; $P<0.001$).

Analysis by indication for catheterization revealed no sex differences in in-laboratory complications and similar or lower rates of bleeding complications among women (Table 4). Unadjusted outcomes for patients with ACS showed that women had lower mortality, all-cause rehospitalization, unplanned PCI, and unplanned CABG. Women and men with ACS had similar rates of rehospitalization for MI and unplanned catheterizations (Table 4). Adjusted models, including the interaction between sex and indication,

showed women with ACS were less likely to die (HR, 0.45; CI, 0.23–0.88; $P=0.014$), but equally likely to be rehospitalized, require an unplanned catheterization, PCI, or CABG (Figure 1). Women with elective catheterizations had lower rates of death, rehospitalization for all causes, and unplanned catheterizations by 1 year (Figure 2). Women and men undergoing catheterization for other indications had no significant differences in outcomes (Figure 3).

Discussion

This is the first study of national VA data to investigate sex differences in the clinical characteristics, treatment, and outcomes of veterans undergoing cardiac catheterization. Our results show that women veterans had fewer classic CAD risk factors than men but higher rates of obesity, depression, and PTSD. Women more often presented with chest pain as an indication for catheterization than men. Furthermore, women veterans had lower rates of obstructive CAD, similar or lower rates of procedural complications, and lower 1-year death and all-cause rehospitalization rates, regardless of catheterization indication. These findings suggest that the clinical presentation that prompts referral of women veterans to VA catheterization laboratories is not as attributable to obstructive CAD as men. Accordingly, exploration of other causes for women's clinical presentation, such as endothelial dysfunction or concurrent psychological comorbidities, should occur.

Our findings both contradict and confirm prior studies of women and cardiovascular disease in several important ways. In contrast to the general population, in which women are typically older and have more cardiovascular risk factors than men,⁷ women veterans were younger and had fewer conventional risk factors. In addition, women veterans had higher rates of depression and PTSD than men veterans, highlighting mental health as a potential risk factor for CAD and adverse outcomes.^{27,28} Depression, anxiety, and other psychological disorders can also manifest physical symptoms, such as chest pain, even in the absence of CAD.^{29–31} Because depression and PTSD are associated with decreased quality of life, persistent chest pain, increased cardiovascular risk, and worse cardiovascular outcomes,^{32–35} the high rates observed in the veteran population clearly warrant attention.

Similar to studies in civilian populations, women veterans have less obstructive CAD than men.^{8,15,36} Women had normal coronary arteries more often than men, regardless of catheterization indication. In the ACS population, women were also more likely to have nonobstructive CAD and less multivessel disease. Possible reasons for less obstructive disease in women with cardiac symptoms have been explored, including disorders of the coronary microvasculature and endothelial dysfunction.^{37–39} Studies have shown that PTSD is associated with chest pain and that there is an association between PTSD and subsequent development of CAD, although this association has not been studied in women veterans specifically. Postulated mechanisms include sympathetic activation, which may over time result in cardiac autonomic dysfunction.^{32,33} However, the low diagnostic yield of cardiac catheterization among women veterans might also indicate that screening and referral for catheterization needs improvement. Additional research is needed to better understand the referral

Table 4. Coronary Anatomy, Postprocedural Medications, and Unadjusted Outcomes by Indication

	ACS			Elective			Other		
	Women (N=401)	Men (N=14 052)	P Value	Women (N=2132)	Men (N=46 309)	P Value	Women (N=648)	Men (N=22 394)	P Value
Coronary anatomy and treatment									
3-Vessel/left main obstructive, % (n)	11.2 (45)	26.8 (3772)	<0.0001	3.7 (78)	15.9 (7357)	<0.0001	2.9 (19)	13.4 (3003)	<0.0001
2-Vessel obstructive, % (n)	13.5 (54)	22.8 (3202)	<0.0001	4.3 (91)	13.4 (6222)	<0.0001	3.9 (25)	10.8 (2418)	<0.0001
1-Vessel obstructive, % (n)	31.2 (125)	32.2 (4527)	0.6591	10.3 (219)	20.8 (9633)	<0.0001	9.7 (63)	17.6 (3943)	<0.0001
Nonobstructive, % (n)	22.7 (91)	11.7 (1645)	<0.0001	29.3 (624)	29.1 (13 473)	0.8622	30.6 (198)	32.9 (7364)	0.2134
Normal, % (n)	19.2 (77)	5 (699)	<0.0001	50 (1065)	18.4 (8534)	<0.0001	49.2 (319)	21.8 (4875)	<0.0001
PCI treatment, % (n)	38.2 (153)	48.1 (6764)	<0.0001	8.9 (190)	20 (9252)	<0.0001	4.5 (29)	9.2 (2054)	<0.0001
Postprocedural medications									
Beta-blocker, % (n)	74.3 (298)	83.3 (11 708)	<0.0001	51.1 (1089)	66.1 (30 612)	<0.0001	58.2 (377)	71.5 (16 017)	<0.0001
Statin, % (n)	74.6 (299)	84.2 (11 838)	<0.0001	56.5 (1204)	70.8 (32 775)	<0.0001	49.4 (320)	63.8 (14 294)	<0.0001
Clopidogrel, prasugrel, % (n)	50.4 (202)	63.6 (8933)	<0.0001	12.6 (269)	29.9 (13 867)	<0.0001	9.6 (62)	16.7 (3740)	<0.0001
Long-acting Nitrate, % (n)	44.9 (180)	49.4 (6940)	0.0755	23.3 (497)	31 (14 358)	<0.0001	14 (91)	20.6 (4604)	<0.0001
Calcium channel blocker, % (n)	21.9 (88)	21.1 (2963)	0.6776	23.3 (496)	24.5 (11 329)	0.2075	23 (149)	20.9 (4682)	0.1983
Procedural complications									
In-laboratory complications, % (n)	2.5 (10)	1.9 (261)	0.3543	1.2 (26)	1.4 (653)	0.4642	2 (13)	1.8 (404)	0.7036
Transfusion within 30 days, % (n)	8 (32)	10.1 (1418)	0.1653	1.7 (37)	4.1 (1914)	<0.0001	6.2 (40)	6.7 (1507)	0.5767
Unadjusted 1-year outcomes									
Death, % (n)	3.2 (13)	9.4 (1320)	<0.0001	1.3 (28)	3.1 (1443)	<0.0001	6.9 (45)	8.5 (1905)	0.1589
Rehospitalization all-cause, % (n)	37.7 (151)	42.5 (5968)	0.0543	24.7 (526)	36.5 (16 883)	<0.0001	45.7 (296)	51.5 (11 532)	0.0035
Rehospitalization for MI (day 15-1 year), % (n)	2.7 (11)	2.6 (359)	0.8138	0.5 (10)	0.9 (397)	0.0548	0.5 (3)	0.9 (196)	0.2635
Repeat catheterization, % (n)	17.2 (69)	18.6 (2615)	0.4764	5.6 (120)	10.8 (5010)	<0.0001	7.7 (50)	9.6 (2151)	0.1068
Unplanned PCI, % (n)	8.5 (34)	11.2 (1580)	0.083	2.1 (45)	4.9 (2257)	<0.0001	2.5 (16)	3.8 (861)	0.0712
Unplanned CABG, % (n)	1.2 (5)	3.8 (527)	0.0087	1.1 (23)	5.1 (2365)	<0.0001	1.9 (12)	6 (1335)	<0.0001

ACS indicates acute coronary syndrome; CABG, coronary artery bypass grafting; MI, myocardial infarction; and PCI, percutaneous coronary intervention.

patterns at the VA; unfortunately, details about the description and nature of chest pain were not available in the current study. In addition, information about stress test results was not available. Efforts are ongoing to capture this important information in VA patients, and future studies will be able to provide insight into catheterization referral patterns. Additional research is also needed to determine why women veterans with cardiac signs or symptoms have less obstructive disease, and what potential treatments can effectively address their symptoms because it is known that women without obstructive CAD frequently have persistent chest pain⁴⁰ associated with increased healthcare costs,⁴¹ rehospitalization,³⁸ and adverse cardiovascular outcomes, including higher rates of MI and mortality.⁴²

Outcomes for women veterans after diagnostic catheterization were similar or better compared with men veterans. Women had lower mortality at 1 year, even when adjusted for age, presence of obstructive disease, and multiple comorbidities. This is in contrast to some prior research that has indicated that women have worse outcomes than men,¹⁷ but studies are conflicting.⁴³ Although our findings seem reassuring, caution is needed when interpreting these results. Because women have significantly less obstructive CAD than men, but similar rates of rehospitalization for MI and unplanned PCI at 1-year, this may indicate that the presence of nonobstructive disease carries a significant healthcare burden.

This study has important clinical implications, not only for practitioners who will care for increasing numbers of women




Figure 1. Adjusted Cox model results for 1-year adverse outcomes for women veterans, relative to men veterans, undergoing catheterization for acute coronary syndrome (ACS). CABG indicates coronary artery bypass grafting; CI, confidence interval; MI, myocardial infarction; and PCI, percutaneous coronary intervention.




Figure 3. Adjusted Cox model results for 1-year adverse outcomes for women veterans, relative to men veterans, undergoing catheterization for other indications. CABG indicates coronary artery bypass grafting; CI, confidence interval; MI, myocardial infarction; and PCI, percutaneous coronary intervention.

veterans at both VA and non-VA facilities, but also for providing insight into the pathophysiology of CAD in women. Efforts are needed to better understand why women experience chest pain in the absence of obstructive CAD.⁴⁰ The optimal method of diagnosis and therapy for patients with chest pain in the absence of obstructive CAD warrants increased attention; greater emphasis on noninvasive testing methods, such as CT coronary angiography, may be indicated. Finally, the interplay between psychological factors and the presence of chest pain merits further research. As the population of women veterans continues to age and develop additional cardiac risk factors, rates of obstructive CAD and associated complications may rise.

This is the first nationwide study of women veterans undergoing cardiac catheterization. VA CART data are an optimal means of capturing this data; however, several limitations deserve consideration. First, although CART data have been shown to be highly accurate,²⁴ misclassification may still occur, and more detailed information regarding

stress testing types and results was not available. In addition, detailed symptom description, biomarker, and EKG data were not available. In the absence of this granular data, we were unable to explore possible predictors of obstructive or nonobstructive CAD. Second, there is potential for overfitting of the Cox models of long-term outcomes, particularly for rehospitalization caused by MI because of lower event rates in this group. Third, the cause of death could not be separated into cardiac-specific mortality because this is not collected in VA data sets. Additional variables, such as aspirin use and left ventricular ejection fraction, were also not available. Finally, the groups of men and women were significantly different in terms of age and clinical demographics. As a result of these large differences, the use of regression adjustment may not have been sufficient to account for the residual confounding effect of age and other clinical differences. Therefore, these findings should be interpreted accordingly.

Conclusions

This is the first study of national VA data to investigate sex differences in the clinical characteristics, treatment, and outcomes of veterans undergoing cardiac catheterization. Our results show that women veterans had fewer classic CAD risk factors than men but higher rates of obesity, depression, and PTSD. Furthermore, women veterans had lower rates of obstructive CAD, similar or lower rates of procedural complications and subsequent rehospitalization, and lower 1-year death rates, regardless of catheterization indication. These findings suggest that the clinical presentation that prompts referral of women veterans to VA catheterization laboratories is not as attributable to obstructive CAD as it is in men. Accordingly, exploration of other causes for women’s clinical presentation, such as endothelial dysfunction or concurrent psychological comorbidities, should occur. Further investigation into the reasons behind these findings and application of those insights into effective treatments could potentially address some of the sex differences noted in nonveteran cardiac populations.




Figure 2. Adjusted Cox model results for 1-year adverse outcomes for women veterans, relative to men veterans, undergoing catheterization for elective indications. CABG indicates coronary artery bypass grafting; CI, confidence interval; MI, myocardial infarction; and PCI, percutaneous coronary intervention.

Disclosures

None.

References

- Murphy SL, Xu J, Kochanek KD. Deaths: final data for 2010. *Natl Vital Stat Rep*. 2013;61:1–117.
- Cowie CC, Rust KF, Ford ES, Eberhardt MS, Byrd-Holt DD, Li C, Williams DE, Gregg EW, Bainbridge KE, Saydah SH, Geiss LS. Full accounting of diabetes and pre-diabetes in the U.S. population in 1988–1994 and 2005–2006. *Diabetes Care*. 2009;32:287–294. doi: 10.2337/dc08-1296.
- Egan BM, Zhao Y, Axon RN. US trends in prevalence, awareness, treatment, and control of hypertension, 1988–2008. *JAMA*. 2010;303:2043–2050. doi: 10.1001/jama.2010.650.
- Lansky AJ, Ng VG, Maehara A, Weisz G, Lerman A, Mintz GS, De Bruyne B, Farhat N, Niess G, Jankovic I, Lazar D, Xu K, Fahy M, Serruys PW, Stone GW. Gender and the extent of coronary atherosclerosis, plaque composition, and clinical outcomes in acute coronary syndromes. *JACC Cardiovasc Imaging*. 2012;5(3 Suppl):S62–S72. doi: 10.1016/j.jcmg.2012.02.003.
- Wong ND, Lopez V, Tang S, Williams GR. Prevalence, treatment, and control of combined hypertension and hypercholesterolemia in the United States. *Am J Cardiol*. 2006;98:204–208. doi: 10.1016/j.amjcard.2006.01.079.
- Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA*. 2010;303:235–241. doi: 10.1001/jama.2009.2014.
- Hochman JS, Tamis JE, Thompson TD, Weaver WD, White HD, Van de Werf F, Aylward P, Topol EJ, Califf RM. Sex, clinical presentation, and outcome in patients with acute coronary syndromes. Global Use of Strategies to Open Occluded Coronary Arteries in Acute Coronary Syndromes IIb Investigators. *N Engl J Med*. 1999;341:226–232. doi: 10.1056/NEJM199907223410402.
- Bairey Merz CN, Shaw LJ, Reis SE, Bittner V, Kelsey SF, Olson M, Johnson BD, Pepine CJ, Mankad S, Sharaf BL, Rogers WJ, Pohost GM, Lerman A, Quyyumi AA, Sopko G; WISE Investigators. Insights from the NHLBI-Sponsored Women's Ischemia Syndrome Evaluation (WISE) Study: Part II: gender differences in presentation, diagnosis, and outcome with regard to gender-based pathophysiology of atherosclerosis and macrovascular and microvascular coronary disease. *J Am Coll Cardiol*. 2006;47(3 Suppl):S21–S29. doi: 10.1016/j.jacc.2004.12.084.
- Shaw LJ, Miller DD, Romeis JC, Kargl D, Younis LT, Chaitman BR. Gender differences in the noninvasive evaluation and management of patients with suspected coronary artery disease. *Ann Intern Med*. 1994;120:559–566.
- Mark DB, Shaw LK, DeLong ER, Califf RM, Pryor DB. Absence of sex bias in the referral of patients for cardiac catheterization. *N Engl J Med*. 1994;330:1101–1106. doi: 10.1056/NEJM199404213301601.
- Roeters van Lennep JE, Zwinderman AH, Roeters van Lennep HW, Westerveld HE, Plokker HW, Voors AA, Brusckhe AV, van der Wall EE. Gender differences in diagnosis and treatment of coronary artery disease from 1981 to 1997. No evidence for the Yentl syndrome. *Eur Heart J*. 2000;21:911–918. doi: 10.1053/ehj.1999.1941.
- Bowker TJ, Turner RM, Wood DA, Roberts TL, Curzen N, Gandhi M, Thompson SG, Fox KM. A national Survey of Acute Myocardial Infarction and Ischaemia (SAMII) in the U.K.: characteristics, management and in-hospital outcome in women compared to men in patients under 70 years. *Eur Heart J*. 2000;21:1458–1463. doi: 10.1053/ehj.2000.2237.
- Blomkalns AL, Chen AY, Hochman JS, Peterson ED, Trynosky K, Diercks DB, Brogan GX Jr, Boden WE, Roe MT, Ohman EM, Gibler WB, Newby LK; CRUSADE Investigators. Gender disparities in the diagnosis and treatment of non-ST-segment elevation acute coronary syndromes: large-scale observations from the CRUSADE (Can Rapid Risk Stratification of Unstable Angina Patients Suppress Adverse Outcomes With Early Implementation of the American College of Cardiology/American Heart Association Guidelines) National Quality Improvement Initiative. *J Am Coll Cardiol*. 2005;45:832–837. doi: 10.1016/j.jacc.2004.11.055.
- Alfredsson J, Stenestrand U, Wallentin L, Swahn E. Gender differences in management and outcome in non-ST-elevation acute coronary syndrome. *Heart*. 2007;93:1357–1362. doi: 10.1136/hrt.2006.102012.
- Hvelplund A, Galatius S, Madsen M, Rasmussen JN, Rasmussen S, Madsen JK, Sand NP, Tilsted HH, Thyssen P, Sindby E, Højbjerg S, Abildstrøm SZ. Women with acute coronary syndrome are less invasively examined and subsequently less treated than men. *Eur Heart J*. 2010;31:684–690. doi: 10.1093/eurheartj/ehp493.
- Kaul P, Tanguay JF, Newby LK, Hochman JS, Westerhout CM, Califf RM, Tricoci P, Gibson CM, Giugliano RP, Harrington RA, Van de Werf F, Armstrong PW. Association between bleeding and mortality among women and men with high-risk acute coronary syndromes: insights from the Early versus Delayed, Provisional Eptifibatide in Acute Coronary Syndromes (EARLY ACS) trial. *Am Heart J*. 2013;166:723–728. doi: 10.1016/j.ahj.2013.07.014.
- Daly C, Clemens F, Lopez Sendon JL, Tavazzi L, Boersma E, Danchin N, Delahaye F, Gitt A, Julian D, Mulcahy D, Ruzyllo W, Thygesen K, Verheugt F, Fox KM; Euro Heart Survey Investigators. Gender differences in the management and clinical outcome of stable angina. *Circulation*. 2006;113:490–498. doi: 10.1161/CIRCULATIONAHA.105.561647.
- Agha Z, Lofgren RP, VanRuiswyk JV, Layde PM. Are patients at Veterans Affairs medical centers sicker? A comparative analysis of health status and medical resource use. *Arch Intern Med*. 2000;160:3252–3257.
- Kazis LE, Ren XS, Lee A, Skinner K, Rogers W, Clark J, Miller DR. Health status in VA patients: results from the Veterans Health Study. *Am J Med. Qual.* 1999;14:28–38.
- Vimalananda VG, Miller DR, Christiansen CL, Wang W, Tremblay P, Fincke BG. Cardiovascular disease risk factors among women veterans at VA medical facilities. *J Gen Intern Med*. 2013;28 Suppl 2:S517–S523. doi: 10.1007/s11606-013-2381-9.
- Frayne SM, Phipps CS, Friedman SA, Berg E, Ananth L, Iqbal S, Hayes PM, Herrera L. *Women Veterans in the Veterans Health Administration. Volume I. Sociodemographic Characteristics and Use of VHA Care*. Washington, DC: Women's Health Evaluation Initiative, Women Veterans Health Strategic Health Care Group, Veterans Health Administration, Department of Veterans Affairs; 2010.
- Maddox TM, Plomondon ME, Petrich M, Tsai TT, Gethoffer H, Noonan G, Gillespie B, Box T, Fihn SD, Jesse RL, Rumsfeld JS. A national clinical quality program for Veterans Affairs catheterization laboratories (from the Veterans Affairs clinical assessment, reporting, and tracking program). *Am J Cardiol*. 2014;114:1750–1757. doi: 10.1016/j.amjcard.2014.08.045.
- Brindis RG, Fitzgerald S, Anderson HV, Shaw RE, Weintraub WS, Williams JF. The American College of Cardiology-National Cardiovascular Data Registry (ACC-NCDR): building a national clinical data repository. *J Am Coll Cardiol*. 2001;37:2240–2245.
- Byrd JB, Vigen R, Plomondon ME, Rumsfeld JS, Box TL, Fihn SD, Maddox TM. Data quality of an electronic health record tool to support VA cardiac catheterization laboratory quality improvement: the VA Clinical Assessment, Reporting, and Tracking System for Cath Labs (CART) program. *Am Heart J*. 2013;165:434–440. doi: 10.1016/j.ahj.2012.12.009.
- Maddox TM, Stanislawski MA, Grunwald GK, Bradley SM, Ho PM, Tsai TT, Patel MR, Sandhu A, Valle J, Magid DJ, Leon B, Bhatt DL, Fihn SD, Rumsfeld JS. Nonobstructive coronary artery disease and risk of myocardial infarction. *JAMA*. 2014;312:1754–1763. doi: 10.1001/jama.2014.14681.
- Bradley SM, Maddox TM, Stanislawski MA, O'Donnell CI, Grunwald GK, Tsai TT, Ho PM, Peterson ED, Rumsfeld JS. Normal coronary rates for elective angiography in the Veterans Affairs Healthcare System: insights from the VA CART program (veterans affairs clinical assessment reporting and tracking). *J Am Coll Cardiol*. 2014;63:417–426. doi: 10.1016/j.jacc.2013.09.055.
- Lichtman JH, Froelicher ES, Blumenthal JA, Carney RM, Doering LV, Frasure-Smith N, Freedland KE, Jaffe AS, Leifheit-Limson EC, Sheps DS, Vaccarino V, Wulsin L; American Heart Association Statistics Committee of the Council on Epidemiology and Prevention and the Council on Cardiovascular and Stroke Nursing. Depression as a risk factor for poor prognosis among patients with acute coronary syndrome: systematic review and recommendations: a scientific statement from the American Heart Association. *Circulation*. 2014;129:1350–1369. doi: 10.1161/CIR.0000000000000019.
- Mehta PK, Wei J, Wenger NK. Ischemic heart disease in women: A focus on risk factors. *Trends Cardiovasc Med*. 2014. pii:S1050–1738(14)00180-7. doi: 10.1016/j.tcm.2014.10.005.
- Christoph M, Christoph A, Dannemann S, Poitz D, Pfluecke C, Strasser RH, Wunderlich C, Koellner V, Ibrahim K. Mental symptoms in patients with cardiac symptoms and normal coronary arteries. *Open Heart*. 2014;1:e000093. doi: 10.1136/openhrt-2014-000093.
- Rasul F, Stansfeld SA, Hart CL, Gillis C, Smith GD. Common mental disorder and physical illness in the Renfrew and Paisley (MIDSPAN) study. *J Psychosom Res*. 2002;53:1163–1170.

31. Ho KY, Kang JY, Yeo B, Ng WL. Non-cardiac, non-oesophageal chest pain: the relevance of psychological factors. *Gut*. 1998;43:105–110.
32. Kubzansky LD, Koenen KC, Spiro A III, Vokonas PS, Sparrow D. Prospective study of posttraumatic stress disorder symptoms and coronary heart disease in the Normative Aging Study. *Arch Gen Psychiatry*. 2007;64:109–116. doi: 10.1001/archpsyc.64.1.109.
33. Tan G, Fink B, Dao TK, Hebert R, Farmer LS, Sanders A, Pastorek N, Gevirtz R. Associations among pain, PTSD, mTBI, and heart rate variability in veterans of Operation Enduring and Iraqi Freedom: a pilot study. *Pain Med*. 2009;10:1237–1245. doi: 10.1111/j.1526-4637.2009.00712.x.
34. Katon W, Hall ML, Russo J, Cormier L, Hollifield M, Vitaliano PP, Beitman BD. Chest pain: relationship of psychiatric illness to coronary arteriographic results. *Am J Med*. 1988;84:1–9.
35. Boscarino JA. A prospective study of PTSD and early-age heart disease mortality among Vietnam veterans: implications for surveillance and prevention. *Psychosom Med*. 2008;70:668–676. doi: 10.1097/PSY.0b013e31817bcaff.
36. Shaw LJ, Shaw RE, Merz CN, Brindis RG, Klein LW, Nallamothu B, Douglas PS, Krone RJ, McKay CR, Block PC, Hewitt K, Weintraub WS, Peterson ED; American College of Cardiology-National Cardiovascular Data Registry Investigators. Impact of ethnicity and gender differences on angiographic coronary artery disease prevalence and in-hospital mortality in the American College of Cardiology-National Cardiovascular Data Registry. *Circulation*. 2008;117:1787–1801. doi: 10.1161/CIRCULATIONAHA.107.726562.
37. Han SH, Bae JH, Holmes DR Jr, Lennon RJ, Eeckhout E, Barsness GW, Rihal CS, Lerman A. Sex differences in atheroma burden and endothelial function in patients with early coronary atherosclerosis. *Eur Heart J*. 2008;29:1359–1369. doi: 10.1093/eurheartj/ehn142.
38. Johnson BD, Shaw LJ, Buchthal SD, Bairey Merz CN, Kim HW, Scott KN, Doyle M, Olson MB, Pepine CJ, den Hollander J, Sharaf B, Rogers WJ, Mankad S, Forder JR, Kelsey SF, Pohost GM; National Institutes of Health-National Heart, Lung, and Blood Institute. Prognosis in women with myocardial ischemia in the absence of obstructive coronary disease: results from the National Institutes of Health-National Heart, Lung, and Blood Institute-Sponsored Women's Ischemia Syndrome Evaluation (WISE). *Circulation*. 2004;109:2993–2999. doi: 10.1161/01.CIR.0000130642.79868.B2.
39. Reis SE, Holubkov R, Conrad Smith AJ, Kelsey SF, Sharaf BL, Reichek N, Rogers WJ, Merz CN, Sopko G, Pepine CJ; WISE Investigators. Coronary microvascular dysfunction is highly prevalent in women with chest pain in the absence of coronary artery disease: results from the NHLBI WISE study. *Am Heart J*. 2001;141:735–741.
40. Jespersen L, Abildstrøm SZ, Hvelplund A, Prescott E. Persistent angina: highly prevalent and associated with long-term anxiety, depression, low physical functioning, and quality of life in stable angina pectoris. *Clin Res Cardiol*. 2013;102:571–581. doi: 10.1007/s00392-013-0568-z.
41. Shaw LJ, Merz CN, Pepine CJ, Reis SE, Bittner V, Kip KE, Kelsey SF, Olson M, Johnson BD, Mankad S, Sharaf BL, Rogers WJ, Pohost GM, Sopko G; Women's Ischemia Syndrome Evaluation (WISE) Investigators. The economic burden of angina in women with suspected ischemic heart disease: results from the National Institutes of Health-National Heart, Lung, and Blood Institute-sponsored Women's Ischemia Syndrome Evaluation. *Circulation*. 2006;114:894–904. doi: 10.1161/CIRCULATIONAHA.105.609990.
42. Johnson BD, Shaw LJ, Pepine CJ, Reis SE, Kelsey SF, Sopko G, Rogers WJ, Mankad S, Sharaf BL, Bittner V, Bairey Merz CN. Persistent chest pain predicts cardiovascular events in women without obstructive coronary artery disease: results from the NIH-NHLBI-sponsored Women's Ischaemia Syndrome Evaluation (WISE) study. *Eur Heart J*. 2006;27:1408–1415. doi: 10.1093/eurheartj/ehl040.
43. Mehili J, Kastrati A, Dirschinger J, Bollwein H, Neumann FJ, Schömig A. Differences in prognostic factors and outcomes between women and men undergoing coronary artery stenting. *JAMA*. 2000;284:1799–1805.

Characteristics and Outcomes of Women Veterans Undergoing Cardiac Catheterization in the Veterans Affairs Healthcare System: Insights from the VA CART Program

Melinda B. Davis, Thomas M. Maddox, Paula Langner, Mary E. Plomondon, John S. Rumsfeld and Claire S. Duvernoy

Circ Cardiovasc Qual Outcomes. 2015;8:S39-S47; originally published online February 24, 2015;

doi: 10.1161/CIRCOUTCOMES.114.001613

Circulation: Cardiovascular Quality and Outcomes is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2015 American Heart Association, Inc. All rights reserved.

Print ISSN: 1941-7705. Online ISSN: 1941-7713

The online version of this article, along with updated information and services, is located on the World Wide Web at:

http://circoutcomes.ahajournals.org/content/8/2_suppl_1/S39

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation: Cardiovascular Quality and Outcomes* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation: Cardiovascular Quality and Outcomes* is online at:
<http://circoutcomes.ahajournals.org/subscriptions/>