

Sex Differences in the Epidemiology of New-Onset In-Hospital Post–Coronary Artery Bypass Graft Surgery Atrial Fibrillation

A Large Multicenter Study

Giovanni Filardo, PhD, MPH; Gorav Ailawadi, MD; Benjamin D. Pollock, MSPH; Briget da Graca, JD, MS; Danielle M. Sass, MPH, CPH; Teresa K. Phan, MS, MS; Debbie E. Montenegro, MSIS; Vinod Thourani, MD; Ralph Damiano, MD

Background—New-onset atrial fibrillation (AF) after coronary artery bypass graft surgery (CABG) is associated with increased morbidity and poorer long-term survival. Although many studies show differences in outcome in women versus men after CABG, little is known about the sex-specific incidence and characteristics of post-CABG AF.

Methods and Results—Overall, 11 236 consecutive patients without preoperative AF underwent isolated CABG from 2002 to 2010 at 4 US academic medical centers and 1 high-volume specialty cardiac hospital. Data routinely collected for the Society of Thoracic Surgeons database were augmented with details on new-onset post-CABG AF events detected via continuous in-hospital ECG/telemetry monitoring. Unadjusted incidence of post-CABG AF was 29.5% (3312/11 236) overall, 30.2% (2485/8214) in men, and 27.4% (827/3022) in women. After adjustment for Society of Thoracic Surgeons–recognized risk factors, women had significantly lower risk for post-CABG AF (odds ratio [95% confidence interval]=0.75 [0.64–0.89]), shorter first, longest, and total duration of AF episodes (mean difference [95% confidence interval]=−2.7 [−4.7 to −0.8] hours; −4.1 [−6.9 to −1.2] hours; −2.4 [−2.5 to −2.3] hours, respectively). At 48 hours, AF-free probabilities were 77% for women and 72% for men ($P<0.001$). Number of episodes ($P=0.18$), operative mortality ($P=0.048$), stroke ($P=0.126$), and discharge in AF ($P=0.234$) did not differ significantly by sex.

Conclusions—These novel data on sex-specific characteristics of new-onset AF after isolated CABG show that women had lower adjusted risk for post-CABG AF and experienced shorter episodes. Investigation of sex-specific impacts on outcomes is needed to identify optimal strategies for prevention and management to ensure all patients achieve the best possible outcomes. (*Circ Cardiovasc Qual Outcomes*. 2016;9:723-730. DOI: 10.1161/CIRCOUTCOMES.116.003023.)

Key Words: atrial fibrillation ■ coronary artery bypass ■ epidemiology ■ morbidity ■ risk factor ■ women

New-onset atrial fibrillation (AF) after coronary artery bypass graft surgery (CABG) is associated with poorer long-term survival^{1,2} and increased morbidity.^{3,4} Because greater age is associated with higher risk of new-onset post-CABG AF,⁵ the trend toward increasing age and risk profiles in the population of patients undergoing CABG⁶ makes understanding this common complication of cardiac surgery a high priority, so that effective means of prevention and management can be identified and developed. Because women undergoing CABG are typically older and have higher risk profiles than their male counterparts,^{7–11} such research may be particularly important to improving outcomes for women, even though some studies suggest that women have lower risk-adjusted odds of developing post-CABG AF.¹² One study investigating the impact of post-CABG AF on long-term

survival found that it affected women more significantly;¹ another identified it as an independent risk factor for in-hospital mortality.¹³

Although previous studies have looked at the incidence of post-CABG AF in men and women—with some showing it to occur more frequently in women (although not to a statistically significant degree)^{14,15} and others showing similar rates¹⁶ or even lower risk for women^{5,12,17,18}—it is not known how patterns of type, timing, duration, or frequency compare between men and women. We collected detailed data on post-CABG AF events (detected via continuous in-hospital ECG/telemetry monitoring) that occurred in patients who underwent isolated CABG surgery at 4 large academic medical centers and 1 high-volume specialty cardiac hospital in the United States during a period of 9 years. Here, we describe and compare the

Received May 19, 2016; accepted September 9, 2016.

From the Office of the Chief Quality Officer, Baylor Scott & White Health, Dallas, TX (G.F., B.D.P., B.d.G., D.M.S., T.K.P., D.E.M.); The Heart Hospital Baylor Plano, TX (G.F.); Division of Thoracic and Cardiovascular Surgery, University of Virginia, Charlottesville (G.A.); Division of Cardiothoracic Surgery, Emory University, Atlanta, GA (V.T.); and Department of Cardiac Surgery, Washington University School of Medicine and Barnes-Jewish Hospital, St Louis, MO (R.D.).

Correspondence to Giovanni Filardo, PhD, MPH, Office of the Chief Quality Officer, Baylor Scott & White Health, 8080 N Central Expressway, Suite 900, Dallas, TX 75206. E-mail giovanfi@baylorhealth.edu

© 2016 American Heart Association, Inc.

Circ Cardiovasc Qual Outcomes is available at <http://circoutcomes.ahajournals.org>

DOI: 10.1161/CIRCOUTCOMES.116.003023

WHAT IS KNOWN

- New-onset AF after CABG is associated with poorer long-term survival and increased morbidity.
- There is limited population-based data available on the sex-specific incidence and characteristics (timing, duration, and frequency) of post-CABG AF episodes.

WHAT THE STUDY ADDS

- This study provides novel sex-specific data on the timing, duration, and frequency of new-onset post-CABG AF (defined as any episode detected via continuous in-hospital ECG/telemetry monitoring and documented by a physician, regardless of duration or need for treatment) from a large population-based multicenter cohort of >11 000 isolated CABG patients and compares these between women and men.
- It shows that, after adjustment for Society of Thoracic Surgeons-recognized risk factors, women had significantly lower risk for post-CABG AF and shorter durations of first and longest AF episodes and total time in AF.
- Further research is needed to investigate the sex-specific impacts of post-CABG AF on outcomes to identify optimal strategies for prevention and management to ensure all patients achieve the best possible outcomes.

epidemiology of post-CABG AF and its characteristics in men versus women, based on these data.

Methods

Study Setting and Design

This large, multicenter observational study included 11 236 consecutive patients (8214 men; 3022 women) without a history of AF who underwent isolated CABG between January 1, 2002, and December 31, 2010, at Baylor University Medical Center (Dallas, TX), The Heart Hospital at Baylor Plano (Plano, TX), Emory University (Atlanta, GA), University of Virginia (Charlottesville, VA), or Washington University (St Louis, MO). Patients were excluded if they had previous AF, previous valve surgery, preoperative endocarditis, or a ventricular assist device.

The study was approved by the institutional review boards of all centers.

Study Data

Data on patient and operative characteristics (Table 1) routinely collected for the Society of Thoracic Surgeons (STS) database were augmented with detailed data on AF events detected via continuous in-hospital ECG/telemetry monitoring.

Exposure and Outcomes of Interest

The exposure of interest was patient's sex.

The primary outcome measures were (1) the adjusted incidence of new-onset in-hospital post-CABG AF, defined as any episode detected via continuous in-hospital ECG/telemetry monitoring and documented by a physician in the chart, regardless of duration or need for treatment and (2) adjusted time (in hours) from end of CABG surgery to first episode of AF.

Table 1. Patient Characteristics by Sex

Characteristics	Men (n=8214, 73.1%)	Women (n=3022, 26.9%)
Study site		
Baylor*	4005 (48.8%)	1421 (47.0%)
Emory	1597 (19.4%)	698 (23.1%)
University of Virginia	1531 (18.6%)	508 (16.8%)
Washington University	1081 (13.2%)	395 (13.1%)
Age†, y	62.5±10.4	64.8±11.1
Body mass index†, kg/m ²	29.3±5.8	30.0±6.9
Race		
White	6732 (82.0%)	2217 (73.4%)
Black	826 (10.1%)	606 (20.1%)
Hispanic	314 (3.8%)	111 (3.7%)
Asian	160 (2.0%)	38 (1.2%)
Other	182 (2.1%)	50 (1.6%)
Diabetes mellitus	3060 (37.3%)	1471 (48.7%)
Renal failure	223 (2.7%)	102 (3.4%)
Creatinine†, mg/dL	1.26±1.18	1.14±1.17
Chronic lung disease	1531 (18.6%)	696 (23.0%)
Systemic hypertension	6843 (83.3%)	2686 (88.9%)
Peripheral vascular disease	1479 (18.0%)	677 (22.4%)
Cerebrovascular disease	1253 (15.3%)	644 (21.3%)
Time from last myocardial infarction to surgery, h		
No MI	4455 (54.2%)	1630 (53.9%)
≤6	69 (0.9%)	23 (0.8%)
>6 but <24	187 (2.3%)	67 (2.2%)
≥24	3503 (42.6%)	1302 (43.1%)
Current smoker	2322 (28.3%)	808 (26.7%)
Congestive heart failure	1722 (21.0%)	798 (26.4%)
Previous PCI	2322 (28.3%)	832 (27.5%)
Previous coronary bypass	418 (5.1%)	125 (4.1%)
Previous valve surgery/procedure	230 (2.8%)	94 (3.1%)
Preoperative angina pectoris	5173 (63.0%)	1931 (63.9%)
Preoperative ejection fraction†, %	48.4±13.4	51.2±13.3
Left main disease	2084 (25.4%)	706 (23.4%)
Operation		
Elective	5067 (61.7%)	1736 (57.4%)
On-pump	5393 (65.7%)	1825 (60.4%)
Preoperative IABP	1133 (13.8%)	392 (13.0%)

*Two Baylor sites participated in the study: one academic medical center and one high-volume specialty cardiac hospital.

†Mean±SD.

IABP indicates intra-aortic balloon pump; and PCI, percutaneous coronary intervention.

Secondary outcomes included duration (in hours) of the first AF episode, duration (in hours) of the longest AF episode, total time (in hours) spent in AF during hospitalization, and total

number of post-CABG AF episodes experienced during hospitalization. Operative deaths (applying the STS definition¹⁹: death during CABG or any in-hospital death during a continuous post-CABG stay or any death within 30 days of post-CABG discharge), stroke, and discharge in unresolved AF were also assessed.

Statistical Analysis

To assess the association between sex and new-onset post-CABG AF, we used a propensity-adjusted generalized estimating equations model approach,^{20–22} whereas a propensity-adjusted generalized estimating equation–like marginal approach using robust sandwich estimates on a Cox proportional hazards model was used to assess the association between sex and time to first AF.^{20,22}

The propensity adjustment accounted for risk factors recognized by the STS¹⁹ (Table 1); these were modeled as independent variables into a multivariable logistic regression model with sex as the outcome. All continuous variables in the propensity model were fitted using restricted cubic splines with 5 knots.^{20,23} Likewise, a restricted cubic spline function with 5 knots was used to fit the propensity score as an independent variable in:

- (1) a generalized estimating equation model with new-onset post-CABG AF as the dependent variable and sex, time of surgery (incremental month starting from January 2002 [month 1] to December 2010 [month 108]), and the sex*time of surgery interaction term as the independent variables. To avoid assuming a linear association of time of operation with the outcome of interest and to avoid the bias inherent

Table 2. Adjusted AF Outcomes by Sex and Year

	2002 (n=803, 7.1%)	2003 (n=1020, 9.1%)	2004 (n=984, 8.8%)	2005 (n=942, 8.4%)	2006 (n=921, 8.2%)	2007 (n=1124, 10.0%)	2008 (n=1997, 17.8%)	2009 (n=1875, 16.7%)	2010 (n=1573, 14.0%)	Total (N=11 236)
Adjusted new-onset AF*										
Male	37 (31–44)	36 (30–43)	34 (37–41)	35 (29–42)	31 (25–39)	34 (28–40)	32 (27–37)	32 (27–37)	30 (25–35)	33 (28–38)
Female	38 (28–48)	33 (24–43)	22 (15–32)	22 (14–33)	25 (16–35)	28 (20–38)	27 (21–35)	28 (21–35)	23 (17–30)	27 (22–32)
Adjusted time to reach 20% AF incidence, h†										
Male	11 (3–19)	14 (4–26)	18 (5–36)	19 (5–40)	16 (4–31)	14 (4–26)	15 (4–28)	16 (4–30)	26 (6–74)	15 (4–28)
Female	17 (5–34)	23 (6–57)	32 (7–179)	36 (7–305)	29 (6–113)	25 (6–69)	27 (6–89)	29 (6–114)	71 (9–437)	26 (6–80)
Adjusted 48-h AF-free (first episode) probabilities (95% CI)										
Male	0.68 (0.50–0.71)	0.71 (0.68–0.73)	0.73 (0.71–0.76)	0.74 (0.72–0.77)	0.73 (0.70–0.76)	0.71 (0.69–0.74)	0.71 (0.69–0.74)	0.72 (0.70–0.75)	0.77 (0.73–0.80)	0.72 (0.71–0.74)
Female	0.73 (0.69–0.77)	0.76 (0.73–0.79)	0.78 (0.075–0.81)	0.79 (0.76–0.81)	0.77 (0.75–0.80)	0.76 (0.74–0.79)	0.77 (0.75–0.80)	0.77 (0.75–0.80)	0.81 (0.79–0.85)	0.77 (0.76–0.79)
Adjusted time to first AF, mean, h‡										
Male	53 (46–60)	54 (48–59)	55 (48–61)	51 (45–57)	48 (41–55)	51 (45–57)	50 (45–56)	48 (42–54)	45 (35–54)	52 (50–54)
Female	54 (45–63)	55 (47–63)	56 (48–64)	52 (45–60)	50 (42–57)	53 (46–60)	52 (45–59)	50 (42–57)	47 (36–57)	54 (50–58)
Adjusted number of AF episodes‡										
Male (all)	1.3 (1.0–1.6)	1.1 (0.9–1.3)	0.8 (0.6–1.1)	0.8 (0.6–1.1)	1.0 (0.7–1.2)	1.0 (0.8–1.3)	1.0 (0.8–1.2)	1.0 (0.7–1.2)	0.8 (0.5–1.2)	0.9 (0.8–1.0)
Male (≥1 episode of AF)	3.5 (2.7–4.2)	3.0 (2.4–3.5)	2.5 (1.8–3.2)	2.6 (1.9–3.2)	3.0 (2.3–3.7)	3.1 (2.5–3.7)	3.1 (2.5–3.7)	2.9 (2.3–3.6)	3.0 (2.0–4.1)	2.9 (2.6–3.1)
Female (all)	1.2 (0.9–1.6)	1.0 (0.7–1.3)	0.8 (0.5–1.0)	0.7 (0.5–1.0)	0.9 (0.6–1.2)	0.9 (0.7–1.2)	0.9 (0.7–1.2)	0.8 (0.6–1.1)	0.7 (0.3–1.1)	0.8 (0.6–0.9)
Female (≥1 episode of AF)	3.6 (2.6–4.6)	3.1 (2.3–4.0)	2.7 (1.8–3.5)	2.8 (2.0–3.6)	3.2 (2.4–4.1)	3.4 (2.6–4.1)	3.4 (2.7–4.1)	3.2 (2.4–4.0)	3.4 (2.2–4.6)	3.1 (2.7–3.6)
Adjusted first AF duration, mean, h‡										
Male	7.7 (4.9–10.6)	8.3 (6.2–10.5)	9.0 (6.4–11.6)	10.3 (7.9–12.7)	11.1 (8.3–13.9)	10.1 (7.8–12.5)	9.6 (7.4–11.8)	10.0 (7.6–12.5)	6.7 (2.9–10.6)	9.0 (8.1–10.1)
Female	5.1 (1.4–8.8)	5.7 (2.6–8.8)	6.3 (3.0–9.5)	7.6 (4.7–10.5)	8.4 (5.3–11.5)	7.4 (4.6–10.1)	6.8 (4.1–9.5)	7.3 (4.3–10.2)	3.4 (0.1–8.3)	6.3 (4.6–7.9)
Adjusted longest AF duration, h‡										
Male	14 (10–18)	13 (10–16)	12 (8–16)	14 (11–18)	17 (13–21)	15 (11–18)	13 (10–17)	14 (10–17)	10 (4–15)	14 (12–15)
Female	8 (3–14)	8 (3–12)	7 (2–12)	10 (5–14)	12 (8–17)	11 (7–15)	10 (6–14)	10 (6–15)	7 (1–13)	10 (7–12)
Adjusted total AF duration, median, h§										
Male	23 (21–24)	21 (19–22)	20 (19–21)	23 (21–24)	27 (24–28)	24 (22–26)	22 (20–23)	21 (20–23)	20 (18–22)	22 (20–24)
Female	20 (19–21)	19 (18–20)	18 (17–19)	20 (19–21)	24 (23–25)	20 (21–22)	20 (19–21)	19 (18–20)	17 (16–19)	19 (18–21)

Time to first AF, duration of first AF, duration of longest AF, and total AF duration were calculated for the 3312 subjects with ≥1 AF episode; number of AF episodes was calculated for all 11 236 study subjects and for the subgroup of 3312 patients who experienced ≥1 AF episode. AF indicates atrial fibrillation.

*Percent (95% CI).

†80th percentile AF-free probability (90th, 75th).

‡95% CI.

§IQR.

with categorization of continuous variables, time of surgery was also modeled as a continuous covariate with a 5-knot restricted cubic spline;^{20,23}

- (2) a Cox proportional hazards model using robust sandwich estimates with time to new-onset post-CABG AF as the dependent variable and the same covariates and covariate structures listed in bullet (1).

Additionally, both models were tested to assess sex differences between sites using the appropriate interaction terms; no interactions among sites were found (models' $P > 0.95$). Multiple imputation of all covariates used in the propensity model was done by Markov Chain Monte Carlo simulation,^{24,25} to account for missing data (ejection fraction: 7.3% missing and left main disease: 0.4% missing).

The same model framework with the appropriate link function was used to assess adjusted sex differences for the other outcomes of interest. Likewise, models were tested for sex differences and site interaction; no interactions among sites were found (all models' $P > 0.5$).

Estimates from all models were used to compute adjusted odds ratios, hazard ratios, mean differences, median differences, confidence intervals (CIs), and adjusted plots and curves. Model findings were also summarized to describe adjusted sex-specific time to reach 20% AF incidence and probability of remaining AF free for the first 48 hours post CABG.

Results

The overall unadjusted incidence of post-CABG AF was 29.5% (3312/11 236); in men, it was 30.2% (2485/8214), and in women, it was 27.4% (827/3022). Table 1 shows the clinical and demographic characteristics of the study population by sex.

The adjusted new-onset AF-related patient outcomes by year and sex are presented in Table 2. Adjusted differences in AF characteristics between women and men are presented in Table 3, whereas Table 4 summarizes differences between the sexes for other adverse outcomes considered, overall and stratified by new-onset post-CABG AF status.

Figure 1 shows the risk-adjusted predicted probability of new-onset post-CABG AF for men and women over time, as well as the risk-adjusted odds ratios (and 95% CIs) for men versus women developing new-onset post-CABG AF in each year. New-onset post-CABG AF significantly decreased from 2002 to 2010 overall ($P = 0.007$) and in both men and women and was lower in women in all years. During the whole study period, women's risk-adjusted odds of developing post-CABG AF were significantly lower than men's (odds ratio [95% CI] for women versus men = 0.75 [0.64–0.89]).

The adjusted mean (95% CI) number of episodes were similar in men (all men: 0.9 [0.8–1.0]; men with ≥ 1 episode: 2.9 [2.6–3.1]) and women (all women: 0.8 [0.6–0.9]; women with ≥ 1 episode: 3.1 [2.7–3.6]).

Overall, on average, during their hospital stay, men had a statistically significantly greater risk for developing new-onset post-CABG AF than women (hazard ratio [95% CI] for men versus women = 1.16 [1.06–1.28]; Figure 2). Results were similar over the years (Figure 2). Adjusted time to reach 20% post-CABG AF incidence was 26 hours in women and 15 hours in men (Table 2), whereas at 48 hours, women had a 77% probability of being AF free compared with 72% for men ($P < 0.001$; Table 2).

Among those who experienced ≥ 1 episode of AF, the adjusted mean (95% CI) duration of the first episode was significantly shorter in women (6.3 [4.4–7.9] hours) than in men (9.0 [8.1–10.1] hours; difference [95% CI]: -3.1 [-5.1 to -1.2] hours). Likewise, in these patients, the adjusted mean (95% CI) duration of the longest AF episode was significantly shorter in women (9 [7–12] hours) than in men (14 [13–15] hours; difference [95% CI]: -4.6 [-7.3 to -2.0] hours). Adjusted median (95% CI) total time in AF was significantly lower in women

Table 3. Adjusted* Comparison of Post-CABG Atrial Fibrillation Characteristics in Women Vs Men by Year

	2002 (n=803, 7.1%)	2003 (n=1020, 9.1%)	2004 (n=984, 8.8%)	2005 (n=942, 8.4%)	2006 (n=921, 8.2%)	2007 (n=1124, 10.0%)	2008 (n=1997, 17.8%)	2009 (n=1875, 16.7%)	2010 (n=1573, 14.0%)	Total (N=11 236)
Adjusted number of AF episodes (all), mean difference (95% CI)†	-0.1 (-0.4 to 0.3)	-0.1 (-0.4 to 0.2)	-0.1 (-0.3 to 0.2)	-0.1 (-0.3 to 0.1)	-0.1 (-0.3 to 0.1)	-0.1 (-0.3 to 0.1)	-0.1 (-0.3 to 0.1)	-0.1 (-0.3 to 0.1)	-0.1 (-0.4 to 0.2)	-0.1 (-0.3 to 0.1)
adjusted P value (all)	$P = 0.18$
Adjusted number of AF episodes (≥ 1 episode AF), mean difference (95% CI)†	0.2 (-0.8 to 1.1)	0.2 (-0.7 to 1.0)	0.2 (-0.5 to 0.9)	0.2 (-0.4 to 0.8)	0.2 (-0.3 to 0.8)	0.3 (-0.3 to 0.8)	0.3 (-0.3 to 0.9)	0.3 (-0.4 to 1.0)	0.3 (-0.5 to 1.2)	0.3 (-0.3 to 0.8)
P value (≥ 1 episode AF)	$P = 0.23$
Adjusted 1st AF duration, mean difference (95% CI)†	-2.6 (-6.3 to 1.0)	-2.7 (-5.8 to 0.4)	-2.7 (-5.7 to -0.1)	-2.7 (-4.9 to -0.5)	-2.7 (-4.7 to -0.8)	-2.8 (-4.7 to -0.8)	-2.8 (-4.9 to -0.6)	-2.8 (-5.3 to -0.3)	-2.8 (-5.8 to 0.2)	-2.7 (-4.7 to -0.8)
P value	$P = 0.02$
Adjusted longest AF duration, mean difference (95% CI)†	-5.5 (-11 to -0.1)	-5.2 (-9.8 to -0.6)	-4.9 (-8.7 to -1.0)	-4.6 (-7.8 to -1.3)	-4.3 (-7.2 to -1.4)	-4.0 (-6.8 to -1.1)	-3.7 (-6.8 to -0.5)	-3.4 (-7.0 to 0.3)	-3.1 (-7.4 to 1.3)	-4.1 (-6.9 to -1.2)
P value	$P = 0.02$
Adjusted total AF duration, median difference (95% CI)‡	-2.7 (-3.0 to -2.4)	-2.4 (-2.5 to -2.1)	-2.3 (-2.5 to -2.2)	-2.4 (-2.7 to -2.0)	-3.0 (-3.3 to -2.7)	-3.1 (-3.4 to -2.8)	-2.5 (-2.7 to -2.4)	-2.4 (-2.6 to -2.2)	-2.5 (-2.8 to -2.3)	-2.4 (-2.5 to -2.3)
P value	$P < 0.0001$

Time to first AF, duration of first AF, duration of longest AF, and total AF duration were calculated for the 3312 subjects with ≥ 1 AF episode; number of AF episodes was calculated for all 11 236 study subjects and for the subgroup of 3312 patients who experienced ≥ 1 AF episode. AF indicates atrial fibrillation; CABG, coronary artery bypass graft; and CI, confidence interval.

*Adjusted analyses used a propensity-adjusted model controlling for recognized STS risk factors (Table 1) in generalized estimating equations models by study site, with an interaction term between sex and month of surgery to allow mean differences to vary over time.

†Mean and (95% CIs).

‡Median (95% CIs).

Table 4. Risk-Adjusted Nonatrial Fibrillation Outcomes by Sex, Overall, and New-Onset Post-CABG AF Status

Outcome	Overall, % (95% CI)	Male (95% CI)	Female (95% CI)	Odds Ratio (95% CI)	P value
All patients (N=11 236)					
Operative mortality					
Unadjusted	1.7 (1.4–1.9)	1.3 (1.1–1.6)	2.5 (2.0–3.1)	1.9 (1.4–2.6)	<0.0001
Adjusted	1.5 (0.9–2.3)	1.3 (0.9–1.8)	2.2 (1.4–3.4)	1.8 (1.0–3.0)	0.048
Stroke					
Unadjusted	1.6 (1.3–1.8)	1.3 (1.0–1.5)	2.3 (1.8–2.9)	1.8 (1.4–2.5)	<0.0001
Adjusted	1.4 (0.8–2.4)	1.3 (0.9–1.7)	1.8 (1.2–2.8)	1.4 (0.8–2.5)	0.126
AF patients only (n=3312)					
Operative mortality					
Unadjusted	3.4 (2.8–4.1)	2.7 (1.8–4.0)	5.8 (3.7–9.0)	2.2 (1.2–4.1)	0.027
Adjusted	3.3 (2.2–5.0)	2.6 (1.7–3.9)	5.4 (3.2–8.9)	2.1 (1.1–4.3)	0.040
Stroke					
Unadjusted	2.5 (2.0–3.1)	2.0 (1.3–3.2)	3.8 (2.2–6.6)	1.9 (0.9–4.0)	0.070
Adjusted	2.2 (1.3–3.7)	1.8 (1.1–2.9)	3.3 (1.7–6.3)	1.9 (0.8–4.4)	0.094
Discharge home in AF					
Unadjusted	0.8 (0.7–1.0)	0.9 (0.7–1.1)	0.7 (0.4–1.0)	0.8 (0.5–1.3)	0.317
Adjusted	0.7 (0.4–1.3)	0.8 (0.5–1.3)	0.5 (0.2–1.2)	0.7 (0.3–1.6)	0.234
Non-AF patients (n=7924)					
Operative mortality					
Unadjusted	0.8 (0.6–1.1)	0.7 (0.4–1.2)	1.3 (0.7–2.4)	1.8 (0.8–3.9)	0.098
Adjusted	0.8 (0.4–1.4)	0.7 (0.4–1.1)	1.0 (0.4–2.1)	1.4 (0.6–3.8)	0.312
Stroke					
Unadjusted	1.1 (0.9–1.4)	1.0 (1.0–1.5)	1.8 (0.0–2.9)	1.8 (0.9–3.7)	0.063
Adjusted	1.0 (0.6–1.6)	0.9 (0.6–1.4)	1.2 (0.6–2.2)	1.3 (0.6–2.8)	0.332

AF indicates atrial fibrillation; CABG, coronary artery bypass graft; and CI, confidence interval.

(19 [18–21] hours) than in men (22 [20–24] hours; difference [95% CI]: -2.4 [-2.5 to -2.3] hours; $P<0.0001$).

Discussion

Ours is the first large (>11 200 patients), multicenter study to assess both the incidence and characteristics of new-onset AF after isolated CABG and to compare these in women versus men. Using detailed clinical data for patients who underwent CABG during a period of 9 years and including data for all episodes of AF detected via continuous in-hospital ECG/telemetry monitoring to provide an enhanced, comprehensive picture of new-onset AF after isolated CABG, our results reveal important differences between the sexes.

The novel sex-specific data presented here show that after robust risk adjustment to account for patients' clinical and nonclinical risk factors, women undergoing isolated CABG are not only less likely to develop new-onset post-CABG AF than their male counterparts but, among those who do suffer this common complication, they experience shorter first and longest episodes of AF, and spend shorter total amounts of time in AF during the hospital stay.

To our knowledge, this is the first study not only to describe the sex-specific characteristics of post-CABG AF but also to examine all of these (timing, duration, and frequency) and incidence simultaneously, giving a complete picture of both the risk for and clinical course of post-CABG AF in women and men.

Our finding of a lower adjusted incidence of post-CABG AF in women is consistent with the results reported from a relatively large study investigating patients who underwent CABG from 1995 to 2009.¹² Although that study used a slightly different AF definition (episodes of any duration during the hospitalization for the index surgery deemed to be clinically significant by the treating physician) from ours, the propensity-matched odds ratio reported for women versus men of 0.76 (95% CI: 0.65–0.90)¹² was similar to our risk-adjusted result of 0.75 (95% CI: 0.64–0.89). These results are also consistent with previous studies that have identified male sex as an independent risk factor for post-CABG AF.^{17,26,27} Although other studies comparing incidence of post-CABG AF between men and women report finding no significant differences, they examined only unadjusted incidence.^{14–16}




Figure 1. Adjusted* predicted probability, odds ratios comparing sex, and 95% confidence intervals of new-onset post-CABG atrial fibrillation according to sex by year. *Society of Thoracic Surgeons factors listed in Table 1, time of surgery (incremental month starting from January 2002 [month 1] to December 2010 [month 108]), and interaction between sex and time of surgery.

With respect to the significantly shorter duration of AF episodes we observed in women versus men, we found no previous studies with which to compare our results. Although 2 small studies have previously reported on the duration of post-CABG episodes,^{28,29} neither reported sex-specific results. Furthermore, they applied different definitions of AF and patient inclusion/exclusion criteria, limiting the value of comparisons to our findings. We similarly found no previous studies comparing frequency or timing of onset of the first episode of AF between men and women. However, the adjusted mean of 52 hours for men and 54 hours for women we observed for time from CABG to first AF episode among patients who developed AF (Table 2) is consistent with previous observations that post-CABG AF onset tends to peak on days 2 to 3 post surgery.^{18,30}

As described elsewhere (article currently under review), the characteristics (including the STS-recognized risk factors) of our study cohort are similar to those reported for isolated CABG patients from 2000 to 2009 in the national STS adult cardiac surgery database.³¹ Given that >95% of the US cardiac surgery programs contribute to the database, this indicates that our results are likely generalizable to the broader population of US CABG patients. The significantly greater age and higher prevalence of comorbidities such as diabetes mellitus, hypertension, and congestive heart failure among the women in our cohort is also typical of the population of CABG patients;^{9,10,32,33} and the borderline statistically significant higher risk-adjusted operative mortality we observed among the women in our study cohort is consistent with the




Figure 2. Adjusted* probability and hazard ratios (with 95% confidence intervals) for freedom from new-onset post-CABG atrial fibrillation (AF), by sex (2002–2010). *Society of Thoracic Surgeons factors listed in Table 1, time of surgery (incremental month starting from January 2002 [month 1] to December 2010 [month 108]), and interaction between sex and time of surgery.

results reported by several previous studies.^{12,34–38} As with any observational study, we cannot rule out the possibility that unknown factors that confound the association between exposure (sex) and outcome (post-CABG AF) exist. However, the propensity-adjusted generalized estimating equations model approach developed to conduct the statistical analysis considered an extensive list of risk factors recognized by the STS in addition to other important variables, providing a rigorous adjustment for potential confounders.

Future research should investigate how the differences in incidence and duration of post-CABG AF observed here between women and men play out in terms of long-term outcomes. Although it is well established that post-CABG AF is associated with poorer long-term survival,^{1,2,39} it is not yet known whether the magnitude of this association is consistent across sex. It is also not known how the characteristics of post-CABG AF affect long-term survival—nor whether such affects (including their magnitudes) are similar across sex. Given the inconsistencies in the existing evidence about whether (and why) women experience worse, equivalent, or better long-term outcomes after CABG,^{40–44} this is a fertile area for research.

Our results showing significant differences in both the risk-adjusted incidence and duration of AF events between women and men provide important groundwork for future investigations of the sex-specific impacts on survival and of the sex-specific effectiveness of strategies for preventing and managing post-CABG AF. Teasing out differences in the impact of post-operative complications and their management on long-term outcomes between men and women could identify opportunities to improve care for both sexes—and help address the current dearth of information on the effectiveness of treatments for cardiovascular disease specifically in women,^{45,46} something which has been called for to achieve health equity.⁴⁷

Sources of Funding

This work was funded by NIH/National Heart Lung Blood Institute (R01HL103683) and in part by the Bradley Family Endowment to

the Baylor Health Care System Foundation, The Baylor Health Care System Cardiovascular Research Committee (Dallas, TX), and the Discovery Foundation (Dallas, TX).

Disclosures

None.

References

1. El-Chami MF, Kilgo P, Thourani V, Lattouf OM, Delurgio DB, Guyton RA, Leon AR, Puskas JD. New-onset atrial fibrillation predicts long-term mortality after coronary artery bypass graft. *J Am Coll Cardiol.* 2010;55:1370–1376. doi: 10.1016/j.jacc.2009.10.058.
2. Filardo G, Hamilton C, Hebler RF Jr, Hamman B, Grayburn P. New-onset postoperative atrial fibrillation after isolated coronary artery bypass graft surgery and long-term survival. *Circ Cardiovasc Qual Outcomes.* 2009;2:164–169. doi: 10.1161/CIRCOUTCOMES.108.816843.
3. Creswell LL, Schuessler RB, Rosenbloom M, Cox JL. Hazards of postoperative atrial arrhythmias. *Ann Thorac Surg.* 1993;56:539–549.
4. Wu ZK, Iivainen T, Pehkonen E, Laurikka J, Zhang S, Tarkka MR. Fibrillation in patients subjected to coronary artery bypass grafting. *J Thorac Cardiovasc Surg.* 2003;126:1477–1482. doi: 10.1016/S0022.
5. Mathew JP, Parks R, Savino JS, Friedman AS, Koch C, Mangano DT, Browner WS. Atrial fibrillation following coronary artery bypass graft surgery: predictors, outcomes, and resource utilization. MultiCenter Study of Perioperative Ischemia Research Group. *JAMA.* 1996;276:300–306.
6. Ferguson TB, Jr., Hammill BG, Peterson ED, DeLong ER, Grover FL. A decade of change—risk profiles and outcomes for isolated coronary artery bypass grafting procedures, 1990–1999: a report from the STS National Database Committee and the Duke Clinical Research Institute. Society of Thoracic Surgeons. *Ann Thorac Surg.* 2002;73:480–489; discussion 489–490.
7. Al-Alao BS, Parissis H, McGovern E, Tolan M, Young VK. Gender influence in isolated coronary artery bypass graft surgery: a propensity match score analysis of early outcomes. *Gen Thorac Cardiovasc Surg.* 2012;60:417–424. doi: 10.1007/s11748-012-0082-7.
8. Sharoni E, Kogan A, Medalion B, Stamler A, Snir E, Porat E. Is gender an independent risk factor for coronary bypass grafting? *Thorac Cardiovasc Surg.* 2009;57:204–208. doi: 10.1055/s-0029-1185367.
9. Saxena A, Dinh D, Smith JA, Shardey G, Reid CM, Newcomb AE. Sex differences in outcomes following isolated coronary artery bypass graft surgery in Australian patients: analysis of the Australasian Society of Cardiac and Thoracic Surgeons cardiac surgery database. *Eur J Cardiothorac Surg.* 2012;41:755–762. doi: 10.1093/ejcts/ezr039.
10. Koch CG, Weng YS, Zhou SX, Savino JS, Mathew JP, Hsu PH, Saidman LJ, Mangano DT; Ischemia Research and Education Foundation; Multicenter

- Study of Perioperative Ischemia Research Group. Prevalence of risk factors, and not gender per se, determines short- and long-term survival after coronary artery bypass surgery. *J Cardiothorac Vasc Anesth*. 2003;17:585–593.
11. Puskas JD, Edwards FH, Pappas PA, O'Brien S, Peterson ED, Kilgo P, Ferguson TB Jr. Off-pump techniques benefit men and women and narrow the disparity in mortality after coronary bypass grafting. *Ann Thorac Surg*. 2007;84:1447–1454; discussion 1454. doi: 10.1016/j.athoracsur.2007.06.104.
 12. Alam M, Lee VV, Elayda MA, Shahzad SA, Yang EY, Nambi V, Jneid H, Pan W, Coulter S, Wilson JM, Ramanathan KB, Ballantyne CM, Virani SS. Association of gender with morbidity and mortality after isolated coronary artery bypass grafting. A propensity score matched analysis. *Int J Cardiol*. 2013;167:180–184. doi: 10.1016/j.ijcard.2011.12.047.
 13. Ergüneş K, Yilik L, Yetkin U, Lafci B, Bayrak S, Ozpak B, Gurbuz A. Early and mid-term outcomes in female patients undergoing isolated conventional coronary surgery. *J Cardiovasc Thorac Res*. 2014;6:105–110. doi: 10.5681/jcvtr.2014.023.
 14. Eifert S, Kilian E, Beiras-Fernandez A, Juchem G, Reichart B, Lamm P. Early and mid term mortality after coronary artery bypass grafting in women depends on the surgical protocol: retrospective analysis of 3441 on- and off-pump coronary artery bypass grafting procedures. *J Cardiothorac Surg*. 2010;5:90. doi: 10.1186/1749-8090-5-90.
 15. Kokkonen L, Järvinen O, Majahalmel S, Virtanen V, Pehkonen E, Mustonen J, Tarkka M. Atrial fibrillation in elderly patients after coronary artery bypass grafting; gender differences in outcome. *Scand Cardiovasc J*. 2005;39:293–298. doi: 10.1080/140117430510035934.
 16. Vaccarino V, Lin ZQ, Kasl SV, Mattera JA, Roumanis SA, Abramson JL, Krumholz HM. Gender differences in recovery after coronary artery bypass surgery. *J Am Coll Cardiol*. 2003;41:307–314.
 17. Aranki SF, Shaw DP, Adams DH, Rizzo RJ, Couper GS, VanderVliet M, Collins JJ Jr, Cohn LH, Burstin HR. Predictors of atrial fibrillation after coronary artery surgery. Current trends and impact on hospital resources. *Circulation*. 1996;94:390–397.
 18. Zaman AG, Archbold RA, Helft G, Paul EA, Curzen NP, Mills PG. Atrial fibrillation after coronary artery bypass surgery: a model for preoperative risk stratification. *Circulation*. 2000;101:1403–1408.
 19. Shahian DM, O'Brien SM, Filardo G, Ferraris VA, Haan CK, Rich JB, Normand SL, DeLong ER, Shewan CM, Dokholyan RS, Peterson ED, Edwards FH, Anderson RP; Society of Thoracic Surgeons Quality Measurement Task Force. The Society of Thoracic Surgeons 2008 cardiac surgery risk models: part 1—coronary artery bypass grafting surgery. *Ann Thorac Surg*. 2009;88(suppl 1):S2–S22. doi: 10.1016/j.athoracsur.2009.05.053.
 20. Harrell FE, Jr. *Regression Modeling Strategies: With Application to Linear Models, Logistic Regression, and Survival Analysis*. New York: Springer-Verlag; 2001.
 21. Hosmer DW, Lemeshow S, Sturdivant RX. *Applied Logistic Regression*. Hoboken, NJ: John Wiley & Sons, Inc; 2013.
 22. Gharibvand L, Liu L. *Analysis of Survival Data with Clustered Events*. SAS Global Forum 2009. <http://support.sas.com/resources/papers/proceedings09/237–2009.pdf>. Accessed August 1, 2016.
 23. Filardo G, Hamilton C, Hamman B, Ng HK, Grayburn P. Categorizing BMI may lead to biased results in studies investigating in-hospital mortality after isolated CABG. *J Clin Epidemiol*. 2007;60:1132–1139. doi: 10.1016/j.jclinepi.2007.01.008.
 24. Yuan YC. Multiple imputation for missing data: concepts and new developments (version 9.0). 2003. <https://support.sas.com/rnd/app/stat/papers/multipleimputation.pdf>. Accessed October 10, 2016.
 25. Schafer JL, Olsen MK. Multiple imputation for multivariate missing-data problems: a data analyst's perspective. *Multivariate Behav Res*. 1998;33:545–571. doi: 10.1207/s15327906mbr3304_5.
 26. Filardo G, Hamilton C, Hamman B, Hebel RF Jr, Grayburn PA. Relation of obesity to atrial fibrillation after isolated coronary artery bypass grafting. *Am J Cardiol*. 2009;103:663–666. doi: 10.1016/j.amjcard.2008.10.032.
 27. Zacharias A, Schwann TA, Riordan CJ, Durham SJ, Shah AS, Habib RH. Obesity and risk of new-onset atrial fibrillation after cardiac surgery. *Circulation*. 2005;112:3247–3255. doi: 10.1161/CIRCULATIONAHA.105.553743.
 28. Tamis-Holland JE, Kowalski M, Rill V, Firoozi K, Steinberg JS. Patterns of atrial fibrillation after coronary artery bypass surgery. *Ann Noninvasive Electrocardiol*. 2006;11:139–144. doi: 10.1111/j.1542-474X.2006.00095.x.
 29. Taylor AD, Groen JG, Thorn SL, Lewis CT, Marshall AJ. New insights into onset mechanisms of atrial fibrillation and flutter after coronary artery bypass graft surgery. *Heart*. 2002;88:499–504.
 30. Place DG, Peragallo RA, Carroll J, Cusimano RJ, Cheng DC. Postoperative atrial fibrillation: a comparison of off-pump coronary artery bypass surgery and conventional coronary artery bypass graft surgery. *J Cardiothorac Vasc Anesth*. 2002;16:144–148.
 31. ElBardissi AW, Aranki SF, Sheng S, O'Brien SM, Greenberg CC, Gammie JS. Trends in isolated coronary artery bypass grafting: an analysis of the Society of Thoracic Surgeons adult cardiac surgery database. *J Thorac Cardiovasc Surg*. 2012;143:273–281. doi: 10.1016/j.jtcvs.2011.10.029.
 32. Ahmed WA, Tully PJ, Knight JL, Baker RA. Female sex as an independent predictor of morbidity and survival after isolated coronary artery bypass grafting. *Ann Thorac Surg*. 2011;92:59–67. doi: 10.1016/j.athoracsur.2011.02.033.
 33. Lehmkuhl E, Kendel F, Gelbrich G, Dunkel A, Oertelt-Prigione S, Babitsch B, Knosalla C, Bairey-Merz N, Hetzer R, Regitz-Zagrosek V. Gender-specific predictors of early mortality after coronary artery bypass graft surgery. *Clin Res Cardiol*. 2012;101:745–751. doi: 10.1007/s00392-012-0454-0.
 34. Bukkapatnam RN, Yeo KK, Li Z, Amsterdam EA. Operative mortality in women and men undergoing coronary artery bypass grafting (from the California Coronary Artery Bypass Grafting Outcomes Reporting Program). *Am J Cardiol*. 2010;105:339–342. doi: 10.1016/j.amjcard.2009.09.035.
 35. Naughton C, Feneck RO, Roxburgh J. Early and late predictors of mortality following on-pump coronary artery bypass graft surgery in the elderly as compared to a younger population. *Eur J Cardiothorac Surg*. 2009;36:621–627. doi: 10.1016/j.ejcts.2009.04.066.
 36. Edwards FH, Carey JS, Grover FL, Bero JW, Hartz RS. Impact of gender on coronary bypass operative mortality. *Ann Thorac Surg*. 1998;66:125–131.
 37. Vaccarino V, Abramson JL, Veledar E, Weintraub WS. Sex differences in hospital mortality after coronary artery bypass surgery: evidence for a higher mortality in younger women. *Circulation*. 2002;105:1176–1181.
 38. Filardo G, Hamman BL, Pollock BD, da Graca B, Sass DM, Phan TK, Edgerton J, Prince SL, Ring WS. Excess short-term mortality in women after isolated coronary artery bypass graft surgery. *Open Heart*. 2016;3:e000386. doi: 10.1136/openhrt-2015-000386.
 39. Villareal RP, Hariharan R, Liu BC, Kar B, Lee VV, Elayda M, Lopez JA, Rasekh A, Wilson JM, Massumi A. Postoperative atrial fibrillation and mortality after coronary artery bypass surgery. *J Am Coll Cardiol*. 2004;43:742–748. doi: 10.1016/j.jacc.2003.11.023.
 40. Jacobs AK, Kelsey SF, Brooks MM, Faxon DP, Chaitman BR, Bittner V, Mock MB, Weiner BH, Dean L, Winston C, Drew L, Sopko G. Better outcome for women compared with men undergoing coronary revascularization: a report from the bypass angioplasty revascularization investigation (BARI). *Circulation*. 1998;98:1279–1285.
 41. Brandrup-Wognsen G, Berggren H, Hartford M, Hjalmarson A, Karlsson T, Herlitz J. Female sex is associated with increased mortality and morbidity early, but not late, after coronary artery bypass grafting. *Eur Heart J*. 1996;17:1426–1431.
 42. Hassan A, Chiasson M, Buth K, Hirsch G. Women have worse long-term outcomes after coronary artery bypass grafting than men. *Can J Cardiol*. 2005;21:757–762.
 43. Guru V, Fremes SE, Austin PC, Blackstone EH, Tu JV. Gender differences in outcomes after hospital discharge from coronary artery bypass grafting. *Circulation*. 2006;113:507–516. doi: 10.1161/CIRCULATIONAHA.105.576652.
 44. den Ruijter HM, Haitjema S, van der Meer MG, van der Harst P, Rouleau JL, Asselbergs FW, van Gilst WH; IMAGINE Investigators. Long-term outcome in men and women after CABG; results from the IMAGINE trial. *Atherosclerosis*. 2015;241:284–288. doi: 10.1016/j.atherosclerosis.2015.02.039.
 45. Hillis LD, Smith PK, Anderson JL, Bittl JA, Bridges CR, Byrne JG, Cigarroa JE, Disesa VJ, Hiratzka LF, Hutter AM Jr, Jessen ME, Keeley EC, Lahey SJ, Lange RA, London MJ, Mack MJ, Patel MR, Puskas JD, Sabik JF, Selnes O, Shahian DM, Trost JC, Winniford MD. 2011 ACCF/AHA guideline for coronary artery bypass graft surgery: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2011;124:e652–e735. doi: 10.1161/CIR.0b013e31823c074e.
 46. Melloni C, Berger JS, Wang TY, Gunes F, Stebbins A, Pieper KS, Dolor RJ, Douglas PS, Mark DB, Newby LK. Representation of women in randomized clinical trials of cardiovascular disease prevention. *Circ Cardiovasc Qual Outcomes*. 2010;3:135–142. doi: 10.1161/CIRCOUTCOMES.110.868307.
 47. Johnson PA, Fitzgerald T, Salganicoff A, Wood SF, Goldstein JM. Sex-specific medical research: why women's health can't wait. 2014. http://www.brighamandwomens.org/Departments_and_Services/womenshealth/ConnorsCenter/Policy/ConnorsReportFINAL.pdf. Accessed March 8, 2016.

Sex Differences in the Epidemiology of New-Onset In-Hospital Post–Coronary Artery Bypass Graft Surgery Atrial Fibrillation: A Large Multicenter Study
Giovanni Filardo, Gorav Ailawadi, Benjamin D. Pollock, Briget da Graca, Danielle M. Sass, Teresa K. Phan, Debbie E. Montenegro, Vinod Thourani and Ralph Damiano

Circ Cardiovasc Qual Outcomes. 2016;9:723-730; originally published online October 18, 2016;

doi: 10.1161/CIRCOUTCOMES.116.003023

Circulation: Cardiovascular Quality and Outcomes is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2016 American Heart Association, Inc. All rights reserved.

Print ISSN: 1941-7705. Online ISSN: 1941-7713

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circoutcomes.ahajournals.org/content/9/6/723>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation: Cardiovascular Quality and Outcomes* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation: Cardiovascular Quality and Outcomes* is online at:
<http://circoutcomes.ahajournals.org//subscriptions/>